

INFUSION SUITE		PROLASTIN-C INFUSION ORDERS	
Durango Infusion Center 270 E 8th Ave Ste N101 Durango, CO 81301		Phone: 970-828-3500	Fax: 970-828-3501
PATIENT INFORMATION - Include Patient Demographics and Insurance Cards			
Name:		DOB:	
MEDICAL INFORMATION			
ICD10:		Patient Height:	
Patient Weight (kg):		Allergies:	
*Weigh patient prior to each infusion			
REQUIRED TESTING			
<input type="checkbox"/> Serum IgA Level: _____			
Additional labs:			
<input type="checkbox"/> Insert IV		<input type="checkbox"/> Access Port/PICC	
PREMEDICATIONS 30 minutes prior to starting			
<input type="checkbox"/> Acetaminophen:		<input type="checkbox"/> 325mg PO X1	<input type="checkbox"/> 500mg PO X1
<input type="checkbox"/> Diphenhydramine:		<input type="checkbox"/> 25mg IV X1	<input type="checkbox"/> 25mg PO X1
<input type="checkbox"/> Solumedrol:		<input type="checkbox"/> 40mg IV X1	<input type="checkbox"/> 100mg IV X1
<input type="checkbox"/> Antihistamine:		<input type="checkbox"/> 650mg PO X1	<input type="checkbox"/> 1000mg PO X1
<input type="checkbox"/> Additional PRN:		<input type="checkbox"/> 50mg IV X1	<input type="checkbox"/> 50mg PO X1
		<input type="checkbox"/> 125mg IV X1	
		<input type="checkbox"/> Cetirizine 10mg PO X1	
		<input type="checkbox"/> Loratadine 10mg PO X1	
PROLASTIN-C ORDERS			
<input type="checkbox"/> Prolastin-C		60mg/kg IV once weekly X _____	
POST INFUSION			
<input type="checkbox"/> Flush IV line with 25mL NS at the same rate of infusion. D/C IV.			
<input type="checkbox"/> Flush IV line with 25 mL NS. Flush port with 10mL NS, Lock port with 5mL Heparin 10-100U/mL and deaccess			
<input type="checkbox"/> Discharge home			
Referring Provider Printed:			
Referring Provider Signature:		Date:	
Referring Provider Phone:		Referring Provider Fax:	
DNG Provider Printed:			
DNG Provider Signature:		Date:	

**Credentials must be included*

Infusion Directions:

- Remove vial and allow to come to room temp before administration
- Pool Prolastin-C to get intended mg/kg body weight dose into an empty sterile intravenous container.
- Discard and document any drug waste
- Infuse at 0.8 ml/kg/min (approximately 15 minutes)
- Infuse using a 5-to-15-micron infusion filter

Dosing calculator: <https://www.prolastin.com/en/hcp/dosing-and-administration>