

INFUSION SUITE		CINQAIR INFUSION ORDERS	
Durango Infusion Center 270 E 8th Ave Ste N101 Durango, CO 81301		Phone: 970-828-3500	Fax: 970-828-3501
PATIENT INFORMATION - Include Patient Demographics and Insurance Cards			
Name:		DOB:	
MEDICAL INFORMATION			
ICD10:		Patient Height:	
Patient Weight (kg):		Allergies:	
*Weigh patient prior to each infusion			
REQUIRED TESTING			
Additional labs:			
<input type="checkbox"/> Insert IV	<input type="checkbox"/> Access Port/PICC		
PREMEDICATIONS 30 minutes prior to starting			
<input type="checkbox"/> Acetaminophen:	<input type="checkbox"/> 325mg PO X1	<input type="checkbox"/> 500mg PO X1	<input type="checkbox"/> 650mg PO X1 <input type="checkbox"/> 1000mg PO X1
<input type="checkbox"/> Diphenhydramine:	<input type="checkbox"/> 25mg IV X1	<input type="checkbox"/> 25mg PO X1	<input type="checkbox"/> 50mg IV X1 <input type="checkbox"/> 50mg PO X1
<input type="checkbox"/> Solumedrol:	<input type="checkbox"/> 40mg IV X1	<input type="checkbox"/> 100mg IV X1	<input type="checkbox"/> 125mg IV X1
<input type="checkbox"/> Antihistamine:	<input type="checkbox"/> Cetirizine 10mg PO X1 <input type="checkbox"/> Loratadine 10mg PO X1		
<input type="checkbox"/> Additional PRN:			
CINQAIR ORDERS			
<input type="checkbox"/> Cinqair	3mg/kg IV every 4 weeks X _____	Total:	_____ mg
POST INFUSION			
<input type="checkbox"/> Flush IV line with 25mL NS at the same rate of infusion. D/C IV. <input type="checkbox"/> Flush IV line with 25 mL NS. Flush port with 10mL NS, Lock port with 5mL Heparin 10-100U/mL and deaccess <input type="checkbox"/> Discharge home			
Referring Provider Printed:		NPI:	
Referring Provider Signature:		Date:	
Referring Provider Phone:		Referring Provider Fax:	
DNG Provider Printed:			
DNG Provider Signature:		Date:	

***Credentials must be included**

Infusion Directions:

- Remove vial and allow to come to room temp before administration
- Withdraw proper volume of Cinqair from the vial(s) based on recommended weight-based dosage.
- Inject Cinqair into 50mL NS, gently invert to mix.
- Discard and document any waste
- Infuse with filter tubing 0.2micron over 20-50-minute period.
- Infusion time may vary depending on the total volume to be infused as based on patient weight.

Dose Calculator: <https://hcp.cinqair.com/dosage-and-administration.aspx>