DURANGO INFUSION

INFUSION SUITE	VYVGART HYTRULO INFUSION ORDER
Durango Infusion Center 270 E 8th Ave Ste N101 Durango, CO 81301	Phone: 970-828-3500 Fax: 970-828-3501

PATIENT INFORMATION - If Outside Referral, Include Patient Demographics and Insurance Cards	
Name:	DOB:
MEDICAL INFORMATION	
ICD10 / Diagnosis:	Height:
Allergies / Hypersensitivities:	Weight (kg):
	*Weigh patient at each visit
REQUIRED CLINICAL DOCUMENTATION	
☐ For Myasthenia Gravis patients: Anti- AChR antibody and Mg-ADL score	
Additional labs:	
☐ Insert subcutaneous access with winged infusion set	

PREMEDICATIONS	30 minutes prior to starting			
☐ Acetaminophen:	☐ 325mg PO X1	☐ 500mg PO X1	☐ 650mg PO X1	☐ 1000mg PO X1
☐ Diphenhydramine:	☐ 25mg IV X1	☐ 25mg PO X1	☐ 50mg IV X1	☐ 50mg PO X1
☐ Solumedrol:	☐ 40mg IV X1	☐ 100mg IV X1	☐ 125mg IV X1	
☐ Antihistamine	Cetirizine 10mg PO X1		☐ Loratadine 10mg PO X1	
☐ Additional PRN:				

VYVGART HYTRULO ORDERS
☐ gMG: Vyvgart Hytrulo 1,008 mg and hyaluronidase 11,200 units - 5.6 ml SQ infusion over 30 to 90 seconds weekly X 4 weeks
☐ CIDP: Vyvgart Hytrulo 1,008 mg and hyaluronidase 11,200 units - 5.6 ml SQ infusion over 30 to 90 seconds weekly X _____

POST INFUSION	
☐ Discharge home after 30 minute observation	
☐ Discharge home without observation	
Signature:	Date:
Provider Name/Credentials:	NPI: Phone Number:
Provider Name/Credentials:	Provider Name/Credentials:
Provider Name/Credentials:	Provider Name/Credentials:
Provider Name/Credentials:	Provider Name/Credentials:

*No Live Vaccines

Infusion Directions: DO NOT ADMINISTER INTRAVENOUSLY.

- Take the vial out of the refrigerator at least 15 minutes before injecting to allow it to reach room temperature.
- Withdraw the entire content of the vial (including overfill) using a polypropylene syringe and an 18G stainless steel transfer needle.
- Remove large air bubbles if present.
- Fill the tubing of the winged infusion set by gently pressing the syringe plunger until the plunger is at 5.6 mL.
- There should be solution at the end of the winged infusion set needle.
- Choose an injection site on the abdomen (at least 2 to 3 inches away from the navel
- Inject subcutaneously into a pinched skin area at an angle of about 45 degrees over 30 to 90 seconds. Rotate injection sites.
- ***For subsequent MG infusions:*** Patient needs to be re-evaluated. Cannot be started sooner than 50 days from previous cycle