



INFUSION SUITE		NUCALA INJECTION ORDERS	
Durango Infusion Center 270 E 8th Ave Ste N101 Durango, CO 81301		Phone: 970-828-3500	Fax: 970-828-3501
PATIENT INFORMATION - Include Patient Demographics and Insurance Cards			
Name:		DOB:	
MEDICAL INFORMATION			
ICD10:		Patient Height:	
Patient Weight (kg):		Allergies:	
*Weigh patient prior to each infusion			
REQUIRED TESTING			
Additional labs:			
PREMEDICATIONS 30 minutes prior to starting			
<input type="checkbox"/> Acetaminophen:	<input type="checkbox"/> 325mg PO X1	<input type="checkbox"/> 500mg PO X1	<input type="checkbox"/> 650mg PO X1 <input type="checkbox"/> 1000mg PO X1
<input type="checkbox"/> Diphenhydramine:	<input type="checkbox"/> 25mg IV X1	<input type="checkbox"/> 25mg PO X1	<input type="checkbox"/> 50mg IV X1 <input type="checkbox"/> 50mg PO X1
<input type="checkbox"/> Methylprednisolone:	<input type="checkbox"/> 40mg IV X1	<input type="checkbox"/> 100mg IV X1	<input type="checkbox"/> 125mg IV X1
<input type="checkbox"/> Antihistamine:	<input type="checkbox"/> Cetirizine 10mg PO X1 <input type="checkbox"/> Loratadine 10mg PO X1		
<input type="checkbox"/> Additional PRN:			
NUCALA ORDERS			
<input type="checkbox"/> Severe Asthma age 12+	100mg subcutaneous every 4 weeks X _____		
<input type="checkbox"/> Severe Asthma age 6-11	40mg subcutaneous every 4 weeks X _____		
<input type="checkbox"/> For EGPA or HES	300mg subcutaneous every 4 weeks X _____		
* 3 separate 100mg injections for EGPA or HES 300mg dose			
POST INJECTION			
<input type="checkbox"/> Discharge home			
Referring Provider Printed:		NPI:	
Referring Provider Signature:		Date:	
Referring Provider Phone:		Referring Provider Fax:	
DNG Provider Printed:			
DNG Provider Signature:		Date:	

**Credentials must be included*

Injection Directions:

- Reconstitute Nucala with 1.2mL Sterile Water for Injection using a 2-3mL syringe and 21G needle
- The reconstituted solution will contain a concentration of 100mg/mL
- Gently swirl the vial for 10 seconds with a circular motion at 15 second intervals until the powder is dissolved
- Remove the reconstituted Nucala using a new 1mL syringe with 21G to 27G 0.5" needle
- Inject in the thigh, abdomen (greater than 2" around navel), or outer area of upper arm