

INFUSION SUITE	SKYRIZI INFUSION ORDERS
-----------------------	--------------------------------

Durango Infusion Center 270 E 8th Ave Ste N101 Durango, CO 81301	Phone: 970-828-3500 Fax: 970-828-3501
---------------------------------------------------------------------	---------------------------------------

PATIENT INFORMATION - If Outside Referral, Include Patient Demographics and Insurance Cards

Name:	DOB:
-------	------

MEDICAL INFORMATION

ICD10 / Diagnosis:	Height:
--------------------	---------

Allergies / Hypersensitivities:	Weight (kg):
---------------------------------	--------------

*Weigh patient at each visit

REQUIRED CLINICAL DOCUMENTATION

<input type="checkbox"/> TB *Initiation	<input type="checkbox"/> ALT, AST, Bilirubin Prior to initiation and then every 4 weeks
-----------------------------------------	-----------------------------------------------------------------------------------------

Additional labs:

<input type="checkbox"/> Insert IV	<input type="checkbox"/> Access Port/PICC
------------------------------------	-------------------------------------------

PREMEDICATIONS	30 minutes prior to starting
-----------------------	------------------------------

<input type="checkbox"/> Acetaminophen:	<input type="checkbox"/> 325mg PO X1	<input type="checkbox"/> 500mg PO X1	<input type="checkbox"/> 650mg PO X1	<input type="checkbox"/> 1000mg PO X1
-----------------------------------------	--------------------------------------	--------------------------------------	--------------------------------------	---------------------------------------

<input type="checkbox"/> Diphenhydramine:	<input type="checkbox"/> 25mg IV X1	<input type="checkbox"/> 25mg PO X1	<input type="checkbox"/> 50mg IV X1	<input type="checkbox"/> 50mg PO X1
-------------------------------------------	-------------------------------------	-------------------------------------	-------------------------------------	-------------------------------------

<input type="checkbox"/> Methylprednisolone:	<input type="checkbox"/> 40mg IV X1	<input type="checkbox"/> 100mg IV X1	<input type="checkbox"/> 125mg IV X1	
----------------------------------------------	-------------------------------------	--------------------------------------	--------------------------------------	--

<input type="checkbox"/> Antihistamine	<input type="checkbox"/> Cetirizine 10mg PO X1	<input type="checkbox"/> Loratadine 10mg PO X1
----------------------------------------	------------------------------------------------	------------------------------------------------

<input type="checkbox"/> Additional PRN:	
------------------------------------------	--

SKYRIZI ORDERS

<input type="checkbox"/> Skyrizi 600 mg IV in _____ mL NS over 1 hour on week 0, week 4, and week 8 for Crohn's and Plaque Psoriasis

<input type="checkbox"/> Skyrizi 1200 mg IV in _____ mL NS over 2 hours on week 0, week 4, and week 8 for Ulcerative Colitis

POST INFUSION

<input type="checkbox"/> Flush IV line with 25mL NS at the same rate of infusion. D/C IV.

<input type="checkbox"/> Flush IV line with 25 mL NS. Flush port with 10mL NS, Lock port with 5mL Heparin 10-100U/mL and deaccess

<input type="checkbox"/> Discharge home

Signature:	Date:
------------	-------

Provider Name/Credentials:	NPI: Provider Phone:
----------------------------	---------------------------------

Provider Name/Credentials:	Provider Name/Credentials:
----------------------------	----------------------------

Provider Name/Credentials:	Provider Name/Credentials:
----------------------------	----------------------------

Provider Name/Credentials:	Provider Name/Credentials:
----------------------------	----------------------------

Vaccinations: No live vaccines 30 days prior to infusing or during Skyrizi treatment

Infusion Directions:

- Withdraw Skyrizi solution from the vial(s) and inject into an intravenous infusion bag or glass bottle:
 - Crohn's / Plaque Psoriasis - 600mg dose in 100 mL, 250 mL, or 500 mL NS
 - Ulcerative Colitis - 1200 mg dose in 250 mL or 500 mL NS
- Do not shake the vial or diluted solution in the infusion bag or glass bottle
- Allow the diluted Skyrizi to come to room temperature prior to administration
- Infuse over at least 1 hour for **600mg dose** and 2 hours for **1200mg dose**