

INFUSION SUITE		COSENTYX INFUSION ORDERS	
Durango Infusion Center 270 E 8th Ave Ste N101 Durango, CO 81301 Phone: 970-828-3500 Fax: 970-828-3501			
PATIENT INFORMATION - If Outside Referral, Include Patient Demographics and Insurance Cards			
Name:		DOB:	
MEDICAL INFORMATION			
ICD10 / Diagnosis:		Height:	
Allergies / Hypersensitivities:		Weight (kg):	
		*Weigh patient at each visit	
REQUIRED CLINICAL DOCUMENTATION			
☐ TB and HepB prior to initiation			
Additional labs:			
☐ Insert IV		Access Port/PICC	
PREMEDICATIONS 30 minutes prior to starting			
☐ Acetaminophen:	☐ 325mg PO X1	☐ 500mg PO X1	☐ 650mg PO X1 ☐ 1000mg PO X1
☐ Diphenhydramine:	☐ 25mg IV X1	☐ 25mg PO X1	☐ 50mg IV X1 ☐ 50mg PO X1
☐ Methylprednisolone:	☐ 40mg IV X1	☐ 100mg IV X1	☐ 125mg IV X1
☐ Antihistamine	☐ Cetirizine 10mg PO X1		☐ Loratadine 10mg PO X1
☐ Additional PRN:			
COSENTYX ORDERS *Loading dose - 6mg/kg and no max dose. Maintenance dose max is 300mg			
☐ Loading: Cosentyx 6mg/kg IV in _____ mL over 30 minutes			
☐ Maintenance: Cosentyx 1.75mg/kg IV in _____ mL over 30 minutes every 4 weeks			
POST INFUSION			
☐ Flush IV line with 50mL NS at the same rate of infusion. D/C IV.			
☐ Flush IV line with 50mL NS. Flush port with 10mL NS, Lock port with 5mL Heparin 10-100U/mL and deaccess			
☐ Discharge home			
Referring Provider Printed:			
Referring Provider Signature:		Date:	NPI:
Referring Provider Phone:		Referring Provider Fax:	
DNG Provider Printed:			
DNG Provider Signature:		Date:	

- Bring vials to room temperature.
- Withdraw the amount of NS from the saline bag equal to amount of medication to be added.
- Gently invert the bag to avoid foaming.
- Infuse over 30 minutes (filter tubing required - 0.2 micrometer). Flush with 50mL NS following infusion.

Body weight at time of dosing	For the loading dose (6 mg/kg) recommended infusion bag	For maintenance dose (1.75 mg/kg) recommended infusion bag
Greater than 52 kg	100 mL	100 mL
Less than or equal to 52 kg	100 mL	50 mL*

Nursing Considerations:

- Not all patients will require a loading dose
- Avoid live vaccines
- May be administered with or without methotrexate for PsA
- Patients at risk for eczematous eruptions ranging from days to months after first dose
- Monitor patients for the development of IBD