

INFUSION SUITE		VYVGART HYTRULO INFUSION ORDER	
Durango Infusion Center 270 E 8th Ave Ste N101 Durango, CO 81301 Phone: 970-828-3500 Fax: 970-828-3501			
PATIENT INFORMATION - If Outside Referral, Include Patient Demographics and Insurance Cards			
Name:		DOB:	
MEDICAL INFORMATION			
ICD10 / Diagnosis:		Height:	
Allergies / Hypersensitivities:		Weight (kg):	
		*Weigh patient at each visit	
REQUIRED CLINICAL DOCUMENTATION			
☐ For Myasthenia Gravis patients: Anti- AChR antibody and Mg-ADL score			
Additional labs:			
☐ Insert subcutaneous access with winged infusion set			
PREMEDICATIONS 30 minutes prior to starting			
☐ Acetaminophen:	☐ 325mg PO X1	☐ 500mg PO X1	☐ 650mg PO X1 ☐ 1000mg PO X1
☐ Diphenhydramine:	☐ 25mg IV X1	☐ 25mg PO X1	☐ 50mg IV X1 ☐ 50mg PO X1
☐ Methylprednisolone:	☐ 40mg IV X1	☐ 100mg IV X1	☐ 125mg IV X1
☐ Antihistamine	☐ Cetirizine 10mg PO X1		☐ Loratadine 10mg PO X1
☐ Additional PRN:			
VYVGART HYTRULO ORDERS			
☐ gMG: Vyvgart Hytrulo 1,008 mg and hyaluronidase 11,200 units - 5.6 ml SQ infusion over 30 to 90 seconds weekly X 4 weeks			
☐ CIDP: Vyvgart Hytrulo 1,008 mg and hyaluronidase 11,200 units - 5.6 ml SQ infusion over 30 to 90 seconds weekly X _____			
POST INFUSION			
☐ Discharge home after 30 minute observation			
☐ Discharge home without observation			
Referring Provider Printed:			
Referring Provider Signature:		Date:	NPI:
Referring Provider Phone:		Referring Provider Fax:	
DNG Provider Printed:			
DNG Provider Signature:		Date:	

***No Live Vaccines**

Infusion Directions:

DO NOT ADMINISTER INTRAVENOUSLY.

- Take the vial out of the refrigerator at least 15 minutes before injecting to allow it to reach room temperature.
- Withdraw the entire content of the vial (including overfill) using a polypropylene syringe and an 18G stainless steel transfer needle.
- Remove large air bubbles if present.
- Fill the tubing of the winged infusion set by gently pressing the syringe plunger until the plunger is at 5.6 mL.
- There should be solution at the end of the winged infusion set needle.
- Choose an injection site on the abdomen (at least 2 to 3 inches away from the navel)
- Inject subcutaneously into a pinched skin area at an angle of about 45 degrees over 30 to 90 seconds. Rotate injection sites.
- ***For subsequent MG infusions:** Patient needs to be re-evaluated. Cannot be started sooner than 50 days from previous cycle